

# Probability of Independent and Dependent Events

CCM2 Unit 6: Probability

# Independent and Dependent Events

- **Independent Events:** two events are said to be independent when one event has no affect on the probability of the other event occurring.
- **Dependent Events:** two events are dependent if the outcome or probability of the first event affects the outcome or probability of the second.

# Independent Events

Suppose a die is rolled and then a coin is tossed.

- Explain why these events are independent.
  - They are independent because the outcome of rolling a die does not affect the outcome of tossing a coin, and vice versa.
- We can construct a table to describe the sample space and probabilities:

	Roll 1	Roll 2	Roll 3	Roll 4	Roll 5	Roll 6
Head						
Tail						

	Roll 1	Roll 2	Roll 3	Roll 4	Roll 5	Roll 6
Head	1,H	2,H	3,H	4,H	5,H	6,H
Tail	1,T	2,T	3,T	4,T	5,T	6,T

- How many outcomes are there for rolling the die?
  - 6 outcomes
- How many outcomes are there for tossing the coin?
  - 2 outcomes
- How many outcomes are there in the sample space of rolling the die and tossing the coin?
  - 12 outcomes

	Roll 1	Roll 2	Roll 3	Roll 4	Roll 5	Roll 6
Head	1,H	2,H	3,H	4,H	5,H	6,H
Tail	1,T	2,T	3,T	4,T	5,T	6,T

- Is there another way to decide how many outcomes are in the sample space?
  - Multiply the number of outcomes in each event together to get the total number of outcomes.
- Let's see if this works for another situation.

A fast food restaurant offers 5 sandwiches and 3 sides. How many different meals of a sandwich and side can you order?

- If our theory holds true, how could we find the number of outcomes in the sample space?
  - $5 \text{ sandwiches} \times 3 \text{ sides} = 15 \text{ meals}$
- Make a table to see if this is correct.

	Sand. 1	Sand. 2	Sand. 3	Sand. 4	Sand. 5
Side 1					
Side 2					
Side 3					

- Were we correct?

# Probabilities of Independent Events

The probability of independent events is the probability of both occurring, denoted by  $P(A \text{ and } B)$  or  $P(A \cap B)$ .

	Roll 1	Roll 2	Roll 3	Roll 4	Roll 5	Roll 6
Head	1,H	2,H	3,H	4,H	5,H	6,H
Tail	1,T	2,T	3,T	4,T	5,T	6,T

Use the table to find the following probabilities:

1. P(rolling a 3)

$$2/12 = 1/6$$

2. P(Tails)

$$6/12 = \frac{1}{2}$$

3. P(rolling a 3 AND getting tails)

$$1/12$$

4. P(rolling an even)

$$6/12 = \frac{1}{2}$$

5. P(heads)

$$6/12 = \frac{1}{2}$$

6. P(rolling an even AND getting heads)

$$3/12 \text{ or } 1/4$$

What do you notice about the answers to 3 and 6?



# Multiplication Rule of Probability

- The probability of two independent events occurring can be found by the following formula:

$$P(A \cap B) = P(A) \times P(B)$$

# Examples

1. At City High School, 30% of students have part-time jobs and 25% of students are on the honor roll. What is the probability that a student chosen at random has a part-time job and is on the honor roll? Write your answer in context.

$$P(\text{PT job and honor roll}) = P(\text{PT job}) \times P(\text{honor roll}) = .30 \times .25 = .075$$

There is a 7.5% probability that a student chosen at random will have a part-time job and be on the honor roll.

2. The following table represents data collected from a grade 12 class in DEF High School.

Plans after High School			
Gender	University	Community College	Total
Males	28	56	84
Females	43	37	80
Total	71	93	164

Suppose 1 student was chosen at **random** from the grade 12 class.

(a) What is the probability that the student is female?  $20/41$  or  $48.8\%$

(b) What is the probability that the student is going to university?  $71/164$   
or

Now suppose 2 people both randomly chose 1 student from the grade 12 class. Assume that it's possible for them to choose the same student.

(c) What is the probability that the first person chooses a student who is female and the second person chooses a student who is going to university?  $355/1681$  or  $21.1\%$

3. Suppose a card is chosen at random from a deck of cards, replaced, and then a second card is chosen.

- Would these events be independent? How do we know?
  - Yes, because the first card is replaced before the second card is drawn.
- What is the probability that both cards are 7s?
  - $P(7) = 4/52$ , so  $P(7 \text{ and } 7) = P(7) \times P(7) = 4/52 \times 4/52 = 1/169$  or .0059.
  - This means that the probability of drawing a 7, replacing the card and then drawing another 7 is 0.59%

# Dependent Events

- Remember, we said earlier that
  - **Dependent Events:** two events are dependent if the outcome or probability of the first event affects the outcome or probability of the second.
- Let's look at some scenarios and determine whether the events are independent or dependent.

# Determine whether the events are independent or dependent:

1. Selecting a marble from a container and selecting a jack from a deck of cards.
  - Independent
2. Rolling a number less than 4 on a die and rolling a number that is even on a second die.
  - Independent
3. Choosing a jack from a deck of cards and choosing another jack, without replacement.
  - Dependent
4. Winning a hockey game and scoring a goal.
  - Dependent

# Probabilities of Dependent Events

- We cannot use the multiplication rule for finding probabilities of dependent events because the one event affects the probability of the other event occurring.
- Instead, we need to think about how the occurrence of one event will effect the sample space of the second event to determine the probability of the second event occurring.
- Then we can multiply the new probabilities.

# Examples

1. Suppose a card is chosen at random from a deck, the card is NOT replaced, and then a second card is chosen from the same deck. What is the probability that both will be 7s?
  - This is similar the earlier example, but these events are dependent? How do we know?
  - How does the first event affect the sample space of the second event?



1. Suppose a card is chosen at random from a deck, the card is NOT replaced, and then a second card is chosen from the same deck. What is the probability that both will be 7s?
  - Let's break down what is going on in this problem:
    - We want the probability that the first card is a 7, or  $P(1^{\text{st}} \text{ is } 7)$ , and the probability that the second card is a 7, or  $P(2^{\text{nd}} \text{ is } 7)$ .
    - $P(1^{\text{st}} \text{ is } 7) = 4/52$  because there are four 7s and 52 cards
    - How is  $P(2^{\text{nd}} \text{ is } 7)$  changed by the first card being a 7?
    - $P(2^{\text{nd}} \text{ is } 7) = 3/51$
    - $P(1^{\text{st}} \text{ is } 7, 2^{\text{nd}} \text{ is } 7) = 4/52 \times 3/51 = 1/221$  or .0045
    - The probability of drawing two sevens without replacement is 0.45%

2. A box contains 5 red marbles and 5 purple marbles. What is the probability of drawing 2 purple marbles and 1 red marble in succession *without replacement*?

- $P(1^{\text{st}} \text{ purple}) = 5/10$
- $P(2^{\text{nd}} \text{ purple}) = 4/9$
- $P(3^{\text{rd}} \text{ red}) = 5/8$
- $P(\text{purple, purple, red}) = 5/10 \times 4/9 \times 5/8 = 5/36$  or .139
- The probability of drawing a purple, a purple, then a red without replacement is 13.9%

3. In Example 2, what is the probability of first drawing all 5 red marbles in succession and then drawing all 5 purple marbles in succession *without replacement*?

- $P(5 \text{ red then } 5 \text{ purple}) = (5/10)(4/9)(3/8)(2/7)(1/6)(5/5)(4/4)(3/3)(2/2)(1/1) = 1/252 \text{ or } .004$
- The probability of drawing 5 red then 5 purple without replacement is 0.4%
- Explain why the last 5 probabilities above were all equivalent to 1.
- This is because there were only purple marbles left, so the probability for drawing a purple marble was 1.